

Towards the physical point hadronic vacuum polarisation from Möbius DWF

Marina Marinković

UNIVERSITY OF
Southampton

with: P. A. Boyle, A. Jüttner, A. Portelli, ...



European Research Council
Established by the European Commission

SHEP *Southampton
High
Energy
Physics*



RBC-UKQCD Collaboration members

UKQCD

Rudy Arthur (Odense)
Peter Boyle (Edinburgh)
Luigi Del Debbio (Edinburgh)
Shane Drury (Southampton)
Jonathan Flynn (Southampton)
Julien Frison (Edinburgh)
Nicolas Garron (Dublin)
Jamie Hudspith (Toronto)
Tadeusz Janowski (Southampton)
Andreas Juettner (Southampton)
Ava Kamseh (Edinburgh)
Richard Kenway (Edinburgh)
Andrew Lytle (TIFR)
Marina Marinkovic (Southampton)
Brian Pendleton (Edinburgh)
Antonin Portelli (Southampton)
Thomas Rae (Mainz)
Chris Sachrajda (Southampton)
Francesco Sanfilippo (Southampton)
Matthew Spraggs (Southampton)
Tobias Tsang (Southampton)

RBC

Ziyuan Bai (Columbia)
Thomas Blum (UConn/RBRC)
Norman Christ (Columbia)
Xu Feng (Columbia)
Tomomi Ishikawa (RBRC)
Taku Izubuchi (RBRC/BNL)
Luchang Jin (Columbia)
Chulwoo Jung (BNL)
Taichi Kawanai (RBRC)
Chris Kelly (RBRC)
Hyung-Jin Kim (BNL)
Christoph Lehner (BNL)
Jasper Lin (Columbia)
Meifeng Lin (BNL)
Robert Mawhinney (Columbia)
Greg McGlynn (Columbia)
David Murphy (Columbia)
Shigemi Ohta (KEK)
Eigo Shintani (Mainz)
Amarjit Soni (BNL)
Sergey Syritsyn (RBRC)
Oliver Witzel (BU)
Hantao Yin (Columbia)
Jianglei Yu (Columbia)
Daiqian Zhang (Columbia)

Hadronic vacuum polarisation

Can be computed in Euclidean space-time [Blum '02]

$$\Pi_{\mu\nu} = a^4 \sum_x e^{iQx} \langle J_\mu^{em}(x) J_\nu^{em}(0) \rangle$$



- $\Pi_{\mu\nu}(Q) = (Q^2 \delta_{\mu\nu} - Q_\mu Q_\nu) \Pi(Q^2)$
- $\hat{\Pi}(Q^2) = \Pi(Q^2) - \Pi(0)$
- $a_\mu^{HLO} = (\frac{\alpha}{\pi})^2 \int_0^\infty dQ^2 f(Q^2) \times \hat{\Pi}(Q^2)$


Systematic uncertainties to be controlled - general

- 1 Simulations at physical m_π
- 2 Controlled continuum limit, FV effects
- 3 Disconnected diagrams [V. Gülpers, Mon, 14.55] [Della Morte et al. '10]
- 4 Obtaining a real world result: charm quark, isospin effects ...

Systematic uncertainties to be controlled - HVP related

- Conventional simulations do not allow access to sufficiently low Fourier momenta
- Integral is dominated in the region where relative errors are enhanced
- Structure of HVP tensor is such that $\Pi(0)$ is not directly accessible
- Systematic uncertainty introduced by extrapolation

Conventional procedure



- $\Pi(Q^2) = \frac{\Pi_{\mu\nu}(Q^2)}{Q_\mu Q_\nu - \delta_{\mu\nu} Q^2}$
- Transverse projection: $Q_\mu = 0$
- Take only diagonal components $\Pi_{\mu\mu}$
- $a_\mu^{HLO} = (\frac{\alpha}{\pi})^2 \int_0^\infty dQ^2 f(Q^2) \times \hat{\Pi}(Q^2)$

Improving the systematics of connected HVP

Several new methods on the market

- R123 procedure ($\Pi(Q^2 = 0)$, utilising twisted BC formalism) [de Divitiis et al '12]
- Padé approximants [Aubin et al '12]
- Dispersive model study [Golterman et al '13]
- Hybrid strategy [Golterman et al '14] [Mon, 14.15, Sess 1D]
- HPQCD time moments [Chakraborty et al '14] [Mon, 15.15, Sess 1D]
- ...

Challenge: Apply the optimal procedure to physical point data

This work: Fitting Padé approximants on the fresh DWF physical point data

inspired by [Aubin et al. '13]

Improving the systematics of connected HVP

Several new methods on the market

- R123 procedure ($\Pi(Q^2 = 0)$, utilising twisted BC formalism) [de Divitiis et al '12]
- Padé approximants [Aubin et al '12]
- Dispersive model study [Golterman et al '13]
- Hybrid strategy [Golterman et al '14] [Mon, 14.15, Sess 1D]
- HPQCD time moments [Chakraborty et al '14] [Mon, 15.15, Sess 1D]
- ...

Challenge: Apply the optimal procedure to physical point data


This work: Fitting Padé approximants on the fresh DWF physical point data

inspired by [Aubin et al. '13]

Previous RBC-UKQCD computation of a_μ^{HLO} [Boyle et al'11]


Non physical m_π , $a^{-1} \approx 1.3, 1.7, 2.3$ GeV

- Local current at source, conserved at sink
- DWF (Möbius scale=1.0), Iwasaki/DSDR gauge action
- Fitting Q^2 -dependence of $\Pi(Q^2)$ up to $Q_C^2 \approx 2.5 - 9$ GeV²



- Strong m_π dependence
- Eliminate the systematics of chiral extrapolation: computing HVP at m_π^{phys}

RBC-UKQCD $N_f = 2 + 1$ Domain Wall ensembles



- a_μ^{HLO} from DWF for non-physical m_π [Boyle et al '11]
- physical point HVP (•) recently measured \rightarrow preliminary results!

a_μ^{HLO} from DWF at physical pion mass

Physical point lattice parameters:


- Möbius DWF, Iwasaki gauge action
 - $48^3 \times 96 \times 24$, $a^{-1} = 1.73 \text{ GeV}$ -measurements underway
 - $64^3 \times 128 \times 12$, $a^{-1} = 2.31 \text{ GeV}$

HVP with Möbius DWF

- Möbius scale =2.0
- Möbius conserved current [see talk by P.Boyle, Mon 6.10p.m., 2.B]
- Local current at source, conserved at sink
- Point source, 12 source positions


Point vs. stochastic source

- Point source, 12 source positions
- Z(2) wall source, 48 source positions
- (one-end trick) [McNeile et al. '06]



Point vs. stochastic source

- Point source, 12 source positions
- Z(2) wall source, 48 source positions
- (one-end trick) [McNeile et al. '06]
- Comparison (12 src. positions each, **log scale on y-axis**)
- Point src. better in low- Q^2 region ($Q^2 < \sim 0.2 \text{ GeV}^2$)




Physical point HVP from $N_f = 2 + 1$ DWF

Physical point data:

- $L/a = 48^3 \times 94 \times 24$, $a^{-1} = 1.73 \text{ GeV}$
- $\Pi(Q^2)$ convergent sequence of PAs [Aubin et al, '13]
 - VMD is unreliable
- Padé approximants $[N,D]$


$$\Pi_{[N,D]}(Q^2) = \frac{\sum_{n=0}^{N-1} a_n Q^{2n}}{1 + \sum_{m=1}^D b_m Q^{2m}}$$

Physical point HVP from $N_f = 2 + 1$ DWF





- $L/a = 48, a^{-1} = 1.73 \text{ GeV}, m_\pi = 138 \text{ MeV}$
- $Q_C^2 = 1.5 \text{ GeV}^2$

Physical point HVP from $N_f = 2 + 1$ DWF



- $L/a = 48, a^{-1} = 1.73 \text{ GeV}, m_\pi = 138 \text{ MeV}$
- $Q_C^2 = 1.5 \text{ GeV}^2$


Physical point HVP from $N_f = 2 + 1$ DWF



- **Left:** Physical point data (Möbius DWF)
- **Right:** Dispersive model study [Golterman et al. '13]
- Same qualitative behaviour - Padé [2,2] looks acceptable
- Nevertheless, even for Padé [2,2]
 - Removing correlations
 - Results for different choice of Q_C^2 not compatible
- Quoting the value for a_μ^{HLO} would be premature

Physical point HVP from $N_f = 2 + 1$ DWF

Light and strange contributions separated



Limited statistics (28 meas. config.) with physical m_π already gives:

• $\frac{\delta a_\mu^{\text{stat.}}}{a_\mu}$ for light contribution is $O(10)$ larger than for strange HVP

Summary and outlook

Summary

- Current status with DWF:
 - physical point data with $\sim 10\%$ stat. errors, measurements underway
 - in addition to the previous non-phys. point computation
- Significant increase signal/noise ratio near $Q^2 = 0$ coming from the light sector
- Large systematics with conventional procedure anticipated


Outlook

- Add another lattice spacing with m_π^{phys}
- Hybrid method [See talks: K.Maltman (Mon, 14.15, 1D)]
- HPQCD time-moment approach [See talks: B.Chakraborty (Mon, 15.15, 1D)] and possible improvements:
 - Discrete moments [See talks: K.Maltman (Mon, 14.15, 1D)]
 - Large volume limit [See talks: C. Lehner (Fri, 15.35, 8D)]
- Ultimate goal: a_μ^{HLO} with full control over syst. and stat. uncertainties ($< 1\%$)

Acknowledgements

- RBC-UKQCD collab. members
- T. Blum, L. Del Debbio, R. J. Hudspith, T. Izubuchi, C. Lehner, R. Lewis, K. Maltman, for useful discussions
- *The research leading to these results has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) ERC grant agreement No 279757*
- The calculations reported here have been done on DIRAC Bluegene/Q computer at the University of Edinburgh's Advanced Computing Facility

Physical point HVP



- $[2, 2]$ Padé fits for different Q_C^2
- Take correlations into account
- Reference $a_\mu^{HLO}(Q_{C \text{ ref}}^2)$ subtracted under bootstrap [$Q_{C \text{ ref}}^2 = 1.5 \text{ GeV}^2$]
- Results for different choice of Q_C^2 not compatible \rightarrow uncontrolled systematics